

Regence

Medical Policy Manual

Medicine, Policy No. 131

Non-Contact Ultrasound Treatments for Wounds

Effective: April 1, 2026

Next Review: February 2027

Last Review: February 2026

IMPORTANT REMINDER

Medical Policies are developed to provide guidance for members and providers regarding coverage in accordance with contract terms. Benefit determinations are based in all cases on the applicable contract language. To the extent there may be any conflict between the Medical Policy and contract language, the contract language takes precedence.

PLEASE NOTE: Contracts exclude from coverage, among other things, services or procedures that are considered investigational or cosmetic. Providers may bill members for services or procedures that are considered investigational or cosmetic. Providers are encouraged to inform members before rendering such services that the members are likely to be financially responsible for the cost of these services.

DESCRIPTION

Non-contact ultrasound treatment devices, which administer low-frequency ultrasound with a saline mist, have been proposed for use in wound healing, including pain management and debridement, via the production, vibration, and movement of micron-sized bubbles in the saline and tissue.

MEDICAL POLICY CRITERIA

Non-contact low-frequency ultrasound is considered **investigational** for the treatment of all wounds.

NOTE: A summary of the supporting rationale for the policy criteria is at the end of the policy.

CROSS REFERENCES

1. [Negative Pressure Wound Therapy in the Outpatient Setting](#), DME, Policy No. 42
2. [Electrostimulation and Electromagnetic Therapy for the Treatment of Wounds](#), DME, Policy No. 83.09
3. [Electromagnetic Therapy Devices](#), Durable Medical Equipment, Policy No. 83.13

BACKGROUND

Ultrasound is defined as a mechanical vibration above the upper threshold of human hearing (greater than 20 KHz) and has been used primarily by physical therapists in the megahertz (MHz) range (1–3 MHz) for the treatment of musculoskeletal disorders. Recently, non-contact ultrasound treatment devices, which administer low-frequency ultrasound in the kilohertz range via a saline mist, have been proposed for use in wound healing, including pain management and debridement. Their proposed mechanism of action is the production, vibration, and movement of micron-sized bubbles in the saline and wound tissue.

REGULATORY STATUS

The following devices were cleared by the US Food and Drug Administration (FDA) through the 510(k) process:

- The MIST Therapy® System, a non-contact ultrasound device, which uses an ultrasound transducer or wand, by Alliqua Biomedical (Langhorne, PA).
- The Qoustic Wound Therapy System™ (formerly called model AR1000) by Arobella Medical LLC. In contrast to MIST system, the Qoustic system uses a contact or non contact techniques to treat wounds. This device is now known as the Qoustic Wound Therapy System™ (K131096).
- Several other devices have been cleared as being substantially equivalent to earlier devices with the FDA product code: NRB.

EVIDENCE SUMMARY

The principal outcomes associated with treatment of wounds, particularly chronic wounds, are complete wound closure (either with or without surgical closure), improvement in the rate or quality of healing (such as the minimization of scarring), treatment of infection, and patient-centered outcomes such as improvements in function or mobility, and minimization of pain.^[1, 2] Outcomes relating to the use of a non-contact low-frequency ultrasound devices for the treatment of wounds are best understood when comparing use of a non-contact low-frequency ultrasound device to a sham device among patients with similar wound type (i.e., burn or chronic diabetic ulcer) receiving standardized wound care regimens. Therefore, data from adequately powered, blinded, randomized sham-controlled trials (RCTs) are required to control for bias and determine whether any treatment effect from non-contact low-frequency ultrasound devices provides a significant advantage over standard wound care.

SYSTEMATIC REVIEWS

Hu (2025) published a network meta-analysis comparing 12 interventions for diabetic foot ulcers, including 99 randomized controlled trials with 7,356 participants.^[3] Surface Under the Cumulative Ranking Curve Stem cell therapy (Surface Under the Cumulative Ranking Curve [SUCRA] = 89.7%; OR = 5.71; 95% CI [2.64-12.34]) and amniotic membrane therapy (SUCRA = 89.2%; OR = 5.11; 95% CI [3.12-8.37]) ranked highest for promoting ulcer healing, while low-frequency ultrasound (OR = 2.20; 95% CI ^[3][0.99-4.91]) and electric stimulation (OR = 1.88; 95% CI [0.87-4.05]) did not demonstrate statistically significant improvements compared to standard of care. Amniotic membrane therapy (SUCRA = 84.7%) significantly reduced healing time by 26.91 days (95% CI [-44.27 to -9.55]), followed by platelet-rich plasma at 21.65 days (95% CI [-33.61 to -9.69]) and negative pressure wound therapy at 16.79 days (95% CI [-31.12 to -2.26]) compared to standard of care. Low-level laser therapy (SUCRA = 93.9%; MD =

34.27; 95% CI [17.35-51.20]) and extracorporeal shockwave therapy (SUCRA = 84.0%; MD = 27.50; 95% CI [11.00-44.00]) demonstrated the greatest improvements in percentage area reduction, while stem cell therapy (SUCRA = 79.9%; OR = 0.12; 95% CI [0.03-0.55]) and hyperbaric oxygen therapy (OR = 0.35; 95% CI [0.16-0.78]) showed the most significant reductions in amputation risk. Study limitations include protocol heterogeneity across included trials and standard of care definitions among studies.

Chen (2023) published a systematic review (SR) evaluating effect of low-frequency ultrasound (LFUS) as an added treatment for chronic wounds.^[4] Studies (n=17) included in the analysis were, RCTs, observational studies, and retrospective studies. The selected studies included 838 subjects with chronic wounds at the baseline of the studies; 412 of them were using the LFUS (225 low-frequency high-intensity contact for diabetic foot wound ulcers, and 187 LFUS low-intensity non-contact for venous leg wound ulcers), and 426 were using standard care (233 sharp débridement for diabetic foot wound ulcers and 193 sham treatments for venous leg wound ulcers). The LFUS high intensity contact for diabetic foot wound ulcers had significantly lower non-healed diabetic foot wound ulcers at ≥ 3 months (OR, 0.37; 95% CI, 0.24-0.56, $p < 0.001$), a higher percentage of diabetic foot wound ulcers area reduction (MD, 17.18; 95% CI, 6.62-27.85, $p = 0.002$) compared with sharp debridement for diabetic foot wound ulcers. The LFUS low-intensity non-contact for a venous leg wound ulcers had a significantly lower non-healed venous leg wound ulcers at ≥ 3 months (OR, 0.31; 95% CI, 0.15-0.62, $p = 0.001$), and higher percentage venous leg wound ulcers area reduction (MD, 18.96; 95% CI, 2.36-35.57, $p = 0.03$) compared with sham treatments for venous leg wound ulcers. The authors conclude that both low-frequency low-intensity non-contact ultrasound and low-frequency high-intensity contact ultrasound adjunctive treatments appear to have a good effect on short-term healing with the endpoints of complete healing and percentage of the wound size decrease. However, they also note that the studies included in this review analysis had low sample size and low numbers of studies within different comparisons.

A Cochrane SR by Cullum (2017) was conducted to evaluate whether venous leg ulcers treated with ultrasound heal more quickly than those not treated with ultrasound.^[5] The review included 11 RCTs that compared either high- or low-frequency ultrasound with non-ultrasound comparator treatments of usual care, sham ultrasound, or alternative leg ulcer treatments. The two trials in the SR that evaluated low-frequency ultrasound^[6, 7] were considered very low-quality due to risk of bias and imprecision. Results from pooled data analyses were unable to determine whether low-frequency ultrasound affects venous ulcer healing at eight and 12 weeks (N=61, RR 3.91, 95% CI 0.47 to 32.85). Consistent with the results of a previous SR,^[8] the authors concluded that it is uncertain whether therapeutic ultrasound (either high- or low-frequency) improves the healing of venous leg ulcers.

Chang (2017) published a SR on the use of low-frequency ultrasound as an adjunct therapy for chronic wound healing.^[9] Of the 25 studies that met selection criteria, four studies were not included due to low quality. The SR did not include meta-analyses due to study heterogeneity and the narrative synthesis did not provide complete information on the range of comparative effects. More than four ultrasound modalities and six wound etiologies were evaluated. Only two studies reported follow-up data, and this follow-up was a short timeframe of 3 months. The authors concluded that while the data supporting the use of low-frequency ultrasound as adjunctive therapy to wound healing is promising, there is not enough high-quality evidence to draw conclusions about its efficacy beyond standard care, a conclusion that is consistent with previous reviews of the relevant literature.^[10, 11]

RANDOMIZED CONTROLLED TRIALS

There have been a number of published unblinded RCTs comparing non-contact low-frequency ultrasound with standard wound care alone in addition to the ones discussed in the SRs above.^[12-18] All of the RCTs used MIST therapy and, other than two trials that did not report a funding source, all were industry funded.^[15, 17] One study addressed diabetic foot ulcers, the population included in the Ennis 2005 RCT, discussed above.^[19] Four RCTs included patients with venous leg ulcers and one RCT evaluated treatment of split thickness graft donor sites. All studies except one, on split thickness graft donor sites, included patients with non-healing wounds; eligibility criteria included wounds that had not healed after at least four weeks. Three studies reported that patients and providers were not blinded but outcome assessment was blinded.^[12-14] The other studies did not mention blinding. Standard care interventions varied somewhat but generally consisted of wound cleaning, noncontact dressings, compression and, if deemed necessary by providers, debridement. Two studies mentioned following national guidelines for the standard care intervention.^[12, 14] Prather did not describe the standard care intervention, and Beheshti reported only that compression was used.^[13, 15] Study characteristics and findings are summarized in Table 1 below.

Table 1. RCTs Comparing Low-Frequency Ultrasound to Standard Care

Study	Initial N Final N	Wounds included	Interventions	Primary Outcome	Outcome Assessment Single-Blinded	Results
Beheshti (2014)	N=90 N=90	Venous leg ulcers (≥ 4 wk)	NLFU: 3 $\times$ /wk until healed (same protocol for HFU) SOC: Compression therapy (visit frequency not reported)	Time to wound healing (months)	NR	NLFU + SOC: 5.70 (SD=1.57) HFU + SOC: 6.10 (SD=1.47) SOC: 8.13 (SD=1.40) 3 groups: p<0.001 HFU vs NLFU: p=0.22
Gibbons (2015)	N=81 N=74	Venous leg ulcers (≥ 30 d)	NLFU: 3 $\times$ /wk for 4 wk SOC: 3 $\times$ /wk for 4 wk	Percent wound area reduction at 4 wk	Yes	NLFU + SOC: - 61% (SD=28.9%) SOC: -45% (SD=32.5%) p=0.002
Olyaie (2013)	N=90 N=90	Venous leg ulcers (≥ 4 wk)	NLFU: 3 $\times$ /wk for 3 mo or until healed (same protocol for HFU) SOC: 3 $\times$ /wk for 3 mo or until healed	Time to wound healing (months)*	NR	NLFU + SOC: 6.65 (SD=1.59) HFU + SOC: 6.86 (SD=2.04) SOC: 8.50 (SD=2.17) 3 groups: p=0.001 HFU vs NLFU: p not reported
Prather (2015)	N=31 N=27	Split-thickness graft donor sites	NLFU: 1 $\times$ /wk for 5 consecutive	Time to wound healing (days)	Yes	NLFU + SOC: 12.1 (SD=6.0)

Study	Initial N Final N	Wounds included	Interventions	Primary Outcome	Outcome Assessment Single-Blinded	Results
			days (after 2-wk run-in period) SOC: 1×/wk for 5 consecutive days (after 2-wk run-in period)			SOC: 21.3 (SD=14.7) p=0.04
White (2015)	N=36 N=36	Venous leg ulcers (≥6 wk)	NLFU: 3×/wk for 8 wk (after 2-wk run-in period) SOC: >1 visit per week for 8 wk	Percent wound area reduction at 13 wk	Yes	NLFU + SOC: -46.6% (SD=38.1%) SOC: -39.2% (SD=38.0%) p=0.565

HFU: high-frequency ultrasound; NLFU: noncontact low-frequency ultrasound; NR: not reported; SOC: standard of care.

* Reported this outcome; did not specify primary outcome.

All but one of the RCTs in Table 1 had statistically significantly better results on the primary outcome with low-frequency ultrasound compared with standard care. However, studies had methodological limitation. In terms of outcome assessment, complete healing is generally considered the most clinically relevant outcome; a 50% reduction in wound area over time and time to heal are also considered to be acceptable outcomes.^[20] A reduction of less than 50%, or wound area reduction without a predefined cutoff is not considered acceptable. The largest number of trials included patients with venous leg ulcers. None of these RCTs had blinded outcome assessment and reported complete healing or one of the other acceptable outcomes as the primary outcome measure. Only one RCT in Table 1, on split thickness graft donor sites, met both of these criteria.^[13] Another limitation of the body of evidence is that some of the standard care interventions involved fewer visits than the NLFU intervention and nonspecific effects of this differential in face-to-face contact could partially explain the difference in findings between intervention and control groups.

NONRANDOMIZED STUDIES

A number of nonrandomized studies described experiences of MIST therapy-treated wound patients.^[21-37] Although these studies contribute to the body of knowledge by providing direction for future research, evidence from these studies does not permit conclusions due to methodological limitations, such as non-random allocation of treatment and lack of appropriate control groups.

PRACTICE GUIDELINE SUMMARY

ASSOCIATION FOR THE ADVANCEMENT OF WOUND CARE

The 2015 update of the Association for the Advancement of Wound Care (AAWC) guideline on treatment of venous ulcers stated that “low-frequency ultrasound may support healing, reduce pain and improve QOL of non-healing venous or mixed etiology venous ulcers” (moderate strength of rating), but also cautions that limited evidence supports enduring benefit or parameters of application.^[38]

In 2010, the Association for the Advancement of Wound Care (AAWC) published a guideline on care of pressure ulcers.^[39] Non-contact ultrasound therapy was included as a potential

second-line intervention if first-line treatments did not result in wound healing, although the strength of the evidence supporting this decision was low (Level C), indicating a lack of sufficient studies on this topic.

AMERICAN COLLEGE OF PHYSICIANS

The American College of Physicians developed a guideline based on evidence on the comparative effectiveness of treatments of pressure ulcers in 2015.^[40] The review of the evidence for therapeutic ultrasound for pressure ulcers concluded that this treatment was similar to controls. Ultrasound is not mentioned in the recommendations.

SOCIETY FOR VASCULAR SURGERY, AMERICAN VENOUS FORUM, AMERICAN PODIATRIC MEDICAL ASSOCIATION

The Society for Vascular Surgery in collaboration with the American Venous Forum published joint guidelines on the management of venous leg ulcers in 2014.^[41] The guidelines recommended adjuvant wound therapy options for venous leg ulcers that fail to demonstrate improvement after four to six weeks of standard wound therapy (strength of recommendation: grade 1; quality of evidence: level B), but suggested against routine ultrasound therapy for venous leg ulcers (strength of recommendation: grade 2; quality of evidence: level B).

The Society for Vascular Surgery in collaboration with the American Podiatric Medical Association published joint guidelines on the management of diabetic foot in 2016.^[42] The guidelines recommended adjuvant therapy that for diabetic foot ulcers that fail to demonstrate more than 50% wound area reduction after four weeks of standard wound therapy. The adjunctive wound therapy options listed in the guidelines are negative pressure therapy, biologics (platelet-derived growth factor, living cellular therapy, extracellular matrix products, amniotic membrane products), and hyperbaric oxygen therapy. Ultrasound therapy is not mentioned as a recommended adjuvant option.

WOUND HEALING SOCIETY

The 2023 Wound Healing Society (WHS) guideline for the treatment of pressure ulcers makes the following recommendations:

"Use of ultrasound may be useful as an adjunct therapy for pressure ulcers unresponsive to standard therapy. Both non-thermal low frequency and high frequency pulsed current ultrasound have been used."^[43]

SUMMARY

There is not enough research to show that non-contact low-frequency ultrasound improves health outcomes for the treatment of wounds. In addition, no practice guidelines recommend non-contact low-frequency ultrasound, for wound healing. Therefore, the use of non-contact low-frequency ultrasound is considered investigational for the treatment of all wounds.

REFERENCES

1. Guidance for industry: chronic cutaneous ulcer and burn wounds-developing products for treatment. *Wound Repair Regen.* 2001;9(4):258-68. PMID: 11679134
2. U.S. Department of Health and Human Services FaDA, Center for Drug Evaluation and Research (CDER), Center for Biologics Evaluation and Research (CBER), Center for Devices and Radiological Health (CDRH) Guidance for Industry Chronic Cutaneous Ulcer and Burn Wounds — Developing Products for Treatment 2006.
3. Hu X, Meng H, Liang J, et al. Comparison of the efficacy of 12 interventions in the treatment of diabetic foot ulcers: a network meta-analysis. *PeerJ.* 2025;13:e19809. PMID: 40821981
4. Chen H, Yu Z, Liu N, et al. The efficacy of low-frequency ultrasound as an added treatment for chronic wounds: A meta-analysis. *International wound journal.* 2023;20(2):448-57. PMID: 35855676
5. Cullum N, Liu Z. Therapeutic ultrasound for venous leg ulcers. *The Cochrane database of systematic reviews.* 2017;5:CD001180. PMID: 28504325
6. Peschen M, Weichenthal M, Schopf E, et al. Low-frequency ultrasound treatment of chronic venous leg ulcers in an outpatient therapy. *Acta dermato-venereologica.* 1997;77(4):311-4. PMID: 9228227
7. Weichenthal M, Mohr P, Stegmann W, et al. Low-frequency ultrasound treatment of chronic venous ulcers. *Wound Repair Regen.* 1997;5:18-22. PMID: 16984453
8. Cullum NA, Al-Kurdi D, Bell-Syer SE. Therapeutic ultrasound for venous leg ulcers. *The Cochrane database of systematic reviews.* 2010(6):CD001180. PMID: 20556749
9. Chang YR, Perry J, Cross K. Low-Frequency Ultrasound Debridement in Chronic Wound Healing: A Systematic Review of Current Evidence. *Plast Surg (Oakv).* 2017;25(1):21-26. PMID: 29026808
10. Tricco AC, Antony J, Vafaei A, et al. Seeking effective interventions to treat complex wounds: an overview of systematic reviews. *BMC Med.* 2015;13:89. PMID: 25899006
11. Voigt J, Wendelken M, Driver V, et al. Low-frequency ultrasound (20-40 kHz) as an adjunctive therapy for chronic wound healing: a systematic review of the literature and meta-analysis of eight randomized controlled trials. *The international journal of lower extremity wounds.* 2011;10(4):190-9. PMID: 22184750
12. Gibbons GW, Orgill DP, Serena TE, et al. A Prospective, Randomized, Controlled Trial Comparing the Effects of Noncontact, Low-frequency Ultrasound to Standard Care in Healing Venous Leg Ulcers. *Ostomy Wound Manage.* 2015;61(1):16-29. PMID: 25581604
13. Prather JL, Tummel EK, Patel AB, et al. Prospective Randomized Controlled Trial Comparing the Effects of Noncontact Low-Frequency Ultrasound with Standard Care in Healing Split-Thickness Donor Sites. *Journal of the American College of Surgeons.* 2015;221(2):309-18. PMID: 25868409
14. White J, Ivins N, Wilkes A, et al. Non-contact low-frequency ultrasound therapy compared with UK standard of care for venous leg ulcers: a single-centre, assessor-blinded, randomised controlled trial. *International wound journal.* 2015. PMID: 25619411
15. Beheshti A, Shafigh Y, Parsa H, et al. Comparison of High-Frequency and MIST Ultrasound Therapy for the Healing of Venous Leg Ulcers. *Advances in clinical and experimental medicine : official organ Wroclaw Medical University.* 2014;23(6):969-75. PMID: 25618125
16. Kavros SJ, Miller JL, Hanna SW. Treatment of ischemic wounds with noncontact, low-frequency ultrasound: the Mayo clinic experience, 2004-2006. *Adv Skin Wound Care.* 2007;20(4):221-6. PMID: 17415030

17. Olyaie M, Rad FS, Elahifar MA, et al. High-frequency and noncontact low-frequency ultrasound therapy for venous leg ulcer treatment: a randomized, controlled study. *Ostomy Wound Manage*. 2013;59(8):14-20. PMID: 23934374
18. Michailidis L, Bergin SM, Haines TP, et al. Healing rates in diabetes-related foot ulcers using low frequency ultrasonic debridement versus non-surgical sharps debridement: a randomised controlled trial. *BMC Res Notes*. 2018;11(1):732. PMID: 30326972
19. Ennis WJ, Foremann P, Mozen N, et al. Ultrasound therapy for recalcitrant diabetic foot ulcers: results of a randomized, double-blind, controlled, multicenter study. *Ostomy Wound Manage*. 2005;51(8):24-39. PMID: 16234574
20. Gottrup F, Apelqvist J, Price P. Outcomes in controlled and comparative studies on non-healing wounds: recommendations to improve the quality of evidence in wound management. *Journal of wound care*. 2010;19(6):237-68. PMID: 20551864
21. Kavros SJ, Liedl DA, Boon AJ, et al. Expedited wound healing with noncontact, low-frequency ultrasound therapy in chronic wounds: a retrospective analysis. *Adv Skin Wound Care*. 2008;21(9):416-23. PMID: 18769168
22. Thomas R. Acoustic pressure wound therapy in the treatment of stage II pressure ulcers. *Ostomy Wound Manage*. 2008;54(11):56-8. PMID: 19037138
23. Bell AL, Cavorsi J. Noncontact ultrasound therapy for adjunctive treatment of nonhealing wounds: retrospective analysis. *Phys Ther*. 2008;88(12):1517-24. PMID: 18849480
24. Cole PS, Quisberg J, Melin MM. Adjuvant use of acoustic pressure wound therapy for treatment of chronic wounds: a retrospective analysis. *J Wound Ostomy Continence Nurs*. 2009;36(2):171-7. PMID: 19287265
25. Medrano S, Beneke MJ. Acoustic pressure wound therapy to debride unstageable pressure ulcers in the acute care setting: a case series. *Ostomy Wound Manage*. 2008;54(12):54-8. PMID: 19104124
26. Caswell D, McNulty BM. Low-frequency, therapeutic ultrasound treatment for congenital ectodermal dysplasia in toddlers. *Ostomy Wound Manage*. 2008;54(10):58-61. PMID: 18927484
27. Howell-Taylor M, Hall MG, Jr., Brownlee lii WJ, et al. Negative pressure wound therapy combined with acoustic pressure wound therapy for infected post surgery wounds: a case series. *Ostomy Wound Manage*. 2008;54(9):49-52. PMID: 18812625
28. Schmuckler J. Acoustic pressure wound therapy to facilitate granulation tissue in sacral pressure ulcers in patients with compromised mobility: a case series. *Ostomy Wound Manage*. 2008;54(8):50-3. PMID: 18716342
29. Serena T, Lee SK, Lam K, et al. The impact of noncontact, nonthermal, low-frequency ultrasound on bacterial counts in experimental and chronic wounds. *Ostomy Wound Manage*. 2009;55(1):22-30. PMID: 19174586
30. Fleming CP. Acoustic pressure wound therapy in the treatment of a vasculopathy-associated digital ulcer: a case study. *Ostomy Wound Manage*. 2008;54(4):62-5. PMID: 18480507
31. Liguori PA, Peters KL, Bowers JM. Combination of negative pressure wound therapy and acoustic pressure wound therapy for treatment of infected surgical wounds: a case series. *Ostomy Wound Manage*. 2008;54(5):50-3. PMID: 18493094
32. Samies J, Gehling M. Acoustic pressure wound therapy for management of mixed partial- and full-thickness burns in a rural wound center. *Ostomy Wound Manage*. 2008;54(3):56-9. PMID: 18456962

33. Serena T. Wound closure and gradual involution of an infantile hemangioma using a noncontact, low-frequency ultrasound therapy. *Ostomy Wound Manage*. 2008;54(2):68-71. PMID: 18401909
34. Waldrop K, Serfass A. Clinical effectiveness of noncontact, low-frequency, nonthermal ultrasound in burn care. *Ostomy Wound Manage*. 2008;54(6):66-9. PMID: 18579927
35. Ennis WJ, Valdes W, Gainer M, et al. Evaluation of clinical effectiveness of MIST ultrasound therapy for the healing of chronic wounds. *Adv Skin Wound Care*. 2006;19(8):437-46. PMID: 17008814
36. Kavros SJ, Schenck EC. Use of noncontact low-frequency ultrasound in the treatment of chronic foot and leg ulcerations: a 51-patient analysis. *J Am Podiatr Med Assoc*. 2007;97(2):95-101. PMID: 17369314
37. Honaker JS, Forston MR, Davis EA, et al. Effects of non contact low-frequency ultrasound on healing of suspected deep tissue injury: a retrospective analysis. *International wound journal*. 2013;10(1):65-72. PMID: 22289135
38. Association for the Advancement of Wound Care (AAWC) Venous Ulcer Guideline. [cited 02/11/2026]. 'Available from:' <https://aawconline.memberclicks.net/assets/appendix%20c%20guideline%20icvug-textformatrecommendations-final%20v42%20changessaved18aug17.pdf>.
39. Association for the Advancement of Wound Care (AAWC). Association for the Advancement of Wound Care guideline of pressure ulcer guidelines. Malvern, Pennsylvania: Association for the Advancement of Wound Care [cited 02/11/2026]. 'Available from:' <https://s3.amazonaws.com/aawc-new/memberclicks/AAWCPressureUlcerGuidelineofGuidelinesAug11.pdf>.
40. Qaseem A, Humphrey LL, Forciea MA, et al. Treatment of pressure ulcers: a clinical practice guideline from the American College of Physicians. *Ann Intern Med*. 2015;162(5):370-9. PMID: 25732279
41. O'Donnell TF, Jr., Passman MA, Marston WA, et al. Management of venous leg ulcers: clinical practice guidelines of the Society for Vascular Surgery (R) and the American Venous Forum. *Journal of vascular surgery*. 2014;60(2 Suppl):3S-59S. PMID: 24974070
42. Hingorani A, LaMuraglia GM, Henke P, et al. The management of diabetic foot: A clinical practice guideline by the Society for Vascular Surgery in collaboration with the American Podiatric Medical Association and the Society for Vascular Medicine. *Journal of vascular surgery*. 2016;63(2 Suppl):3S-21S. PMID: 26804367
43. Gould LJ, Alderden J, Aslam R, et al. WHS guidelines for the treatment of pressure ulcers-2023 update. *Wound Repair Regen*. 2024;32(1):6-33. PMID: 37970711

CODES

Codes	Number	Description
CPT	97610	Low frequency, non-contact, non-thermal ultrasound, including topical application(s), when performed, wound assessment, and instruction(s) for ongoing care, per day
HCPCS	None	

Date of Origin: April 2008