



## Genetic Testing for Myeloid Neoplasms and Leukemia

Effective: June 1, 2025

Next Review: February 2026

Last Review: April 2025

### IMPORTANT REMINDER

Medical Policies are developed to provide guidance for members and providers regarding coverage in accordance with contract terms. Benefit determinations are based in all cases on the applicable contract language. To the extent there may be any conflict between the Medical Policy and contract language, the contract language takes precedence.

PLEASE NOTE: Contracts exclude from coverage, among other things, services or procedures that are considered investigational or cosmetic. Providers may bill members for services or procedures that are considered investigational or cosmetic. Providers are encouraged to inform members before rendering such services that the members are likely to be financially responsible for the cost of these services.

### DESCRIPTION

Genetic testing, including testing for BCR/ABL1 (t(9;22)) translocations and for *ABL1*, *ASXL1*, *CALR*, *CEBPA*, *FLT3*, *IDH1*, *IDH2*, *JAK2*, *KIT*, *MPL*, *NPM1*, *RUNX1*, and/or *TP53* variants may inform the diagnostic, prognostic, and treatment selection processes for myelodysplastic-myeloproliferative neoplasms and select myeloid neoplasms.

### MEDICAL POLICY CRITERIA

**Note:** Please refer to the Cross References section below for genetic testing not addressed in this policy, including but not limited to single-gene testing.

- I. Genetic testing, including panel testing, for *BCR/ABL1* translocation (Philadelphia chromosome) and/or variants in any of the following genes may be considered **medically necessary** for evaluation, diagnosis, and/or treatment monitoring in myeloid neoplasms and leukemia: *JAK2*, *CALR*, *MPL*, *ASXL1*, *IDH1*, *IDH2*, *TP53*, *CEBPA*, *FLT3*, *KIT*, *NPM1* and/or *RUNX1*.
- II. Targeted genetic panel testing for myeloproliferative neoplasms (MPN), myelodysplastic syndromes (MDS), and myelodysplastic myeloproliferative neoplasms (MPN/MDS), including acute myeloid leukemia (AML), may be considered **medically**

**necessary** for patients being evaluated for these disorders (see Policy Guidelines and Table 1).

- III. Genetic testing for *ABL1* may be considered **medically necessary** to evaluate patients when either of the following are met:
  - A. In patients with chronic myelogenous (myeloid) leukemia (CML), to monitor response to tyrosine kinase inhibitor therapy; or
  - B. In patients with Philadelphia chromosome-positive acute lymphoblastic leukemia (Ph+ ALL), to evaluate for tyrosine kinase inhibitor resistance.
- IV. Genetic testing for *ABL1* is considered **investigational** when Criterion III. is not met.
- V. Non-targeted profiling panels for hematologic disorders are considered **investigational** (see Policy Guidelines).

*NOTE: A summary of the supporting rationale for the policy criteria is at the end of the policy.*

## POLICY GUIDELINES

### PANEL TESTING

#### Targeted Panels for Myeloid Neoplasms

Targeted panel testing for myeloid neoplasms, (i.e., MPN, MDS, MPN/MDS, and AML, see Table 1 below) includes panels that are specifically designed to assess variants in patients suspected of having a myeloproliferative neoplasm, a myelodysplastic syndrome, or a disorder with overlapping features. They are generally less than 50 genes and may include the following genes: *ASXL1*, *CALR*, *CBL*, *EZH2*, *KIT*, *FLT3*, *JAK2*, *MPL*, *NMP1*, *CEBPA*, *IDH1*, *IDH2*, and *TP53*.

Examples of targeted panels for MPN/MDS/AML include, but are not limited to:

- NeoTYPE™ Myeloid Disorders Profile (Neogenomics)
- NGS Myeloid 37 Gene panel (Cellnetix)
- MyeloSeq™ (Washington University School of Medicine)
- NGS\_AML Panel (Cellnetix)
- AML Mutation Analysis Panel (Molecular Pathology Laboratory Network)
- Onkosight™ Myeloid Malignancies Panel, MPN Panel, MDS Panel, or AML Panel
- Myeloid MPN/MDS/CMML Comprehensive Panel (Providence)
- Myeloid Gene Panel by NGS (University of Washington)
- TruSight® Myeloid Sequencing Panel

#### Non-targeted Panels

Some commercially available panels are not targeted toward genes that have clinical significance for a specific type of hematolymphoid disorder. They often include testing for a large number of genes that do not have demonstrated clinical utility, as well as testing for many disorders that could be distinguished based on clinical presentation.

Non-targeted panels for hematologic disorders include, but are not limited to:

- FoundationOne Heme (Foundation Medicine)
- FusionPlex Pan-Heme Panel (Laboratory for Precision Diagnostics, University of Washington)
- GeneTrails® Hematologic Malignancies 220 Gene Panel (Knight Diagnostic Laboratories)
- GeneTrails® Heme Fusion Gene Panel
- MyAML® 194 Targeted NGS Gene Panel (Invivoscribe)
- HopeSeq HemeComplete (City of Hope)
- NGS Hematology Molecular Profile (Sonora Quest Laboratories)
- Rapid Heme Panel (Dana-Farber Cancer Institute)
- Hematologic Malignancy Sequencing Panel (Penn Medicine)
- Neo Comprehensive™ - Myeloid Disorders (Neogenomics)

**Table 1. Selected Diagnoses from the World Health Organization Classification of Hematolymphoid Disorders<sup>[1 2]</sup>**

<b>Myeloproliferative neoplasms (MPN)</b>
Chronic myeloid leukemia (CML), <i>BCR-ABL1</i> <sup>+</sup>
Chronic neutrophilic leukemia
Chronic eosinophilic leukemia
Polycythemia vera
Essential thrombocythemia
Primary myelofibrosis
Juvenile myelomonocytic leukemia (JMML)
<b>Mastocytosis</b>
Cutaneous mastocytosis
Systemic mastocytosis
Mast cell sarcoma
<b>Myelodysplastic neoplasms (MDS)</b>
MDS with low blasts and 5q deletion
MDS with low blasts and <i>SF3B1</i> mutation
Myelodysplastic neoplasm with increased blasts
Refractory cytopenia of childhood
Chronic myelomonocytic leukemia (CMML)
<b>Acute myeloid leukemia (AML) and related neoplasms</b>
AML with defining genetic abnormalities
AML, defined by differentiation
Acute basophilic leukemia
Pure erythroid leukemia
Acute megakaryoblastic leukemia
Myeloid sarcoma

## LIST OF INFORMATION NEEDED FOR REVIEW

### REQUIRED DOCUMENTATION:

The information below **must** be submitted for review to determine whether policy criteria are met. If any of these items are not submitted, it could impact our review and decision outcome:

1. Name of the genetic test(s) or panel test
2. Name of the performing laboratory and/or genetic testing organization (more than one may be listed)

3. The exact gene(s) and/or variant(s) being tested
4. Relevant billing codes
5. Brief description of how the genetic test results will guide clinical decisions that would not otherwise be made in the absence testing
6. Medical records related to this genetic test:
  - History and physical exam including any relevant diagnoses related to the genetic testing
  - Sample collection (e.g., blood draw) date
  - Conventional testing and results

## CROSS REFERENCES

1. [Genetic Testing for Hereditary Breast and Ovarian Cancer and Li-Fraumeni Syndrome](#), Genetic Testing, Policy No. 02
2. [Genetic Testing for  \$\alpha\$ -Thalassemia](#), Genetic Testing, Policy No. 19
3. [Genetic and Molecular Diagnostic Testing](#), Genetic Testing, Policy No. 20
4. [ClonoSEQ® Testing for the Assessment of Measurable Residual Disease \(MRD\)](#), Genetic Testing, Policy No. 88
5. [Hematopoietic Cell Transplantation for Acute Myeloid Leukemia](#), Transplant, Policy No. 45.28
6. [Hematopoietic Cell Transplantation for Chronic Myelogenous Leukemia](#), Transplant, Policy No. 45.31
7. [Hematopoietic Cell Transplantation for Acute Lymphoblastic Leukemia](#), Transplant, Policy No. 45.36
8. [Medication Policy Manual](#), Note: Click the link for the appropriate Medication Policy. Once the medication policy site is open, do a find (Ctrl+F) and enter drug name in the find bar to locate the appropriate policy.

## BACKGROUND

### DIAGNOSING MYELOID NEOPLASMS AND ACUTE LEUKEMIA

Myeloid neoplasms may be acute or chronic, are a type of hematologic malignancy, and usually derive from bone marrow progenitor cells that normally develop into erythrocytes, granulocytes (neutrophils, basophils, and eosinophils), monocytes, or megakaryocytes. Classification of myeloid neoplasms and acute leukemias has evolved over the past decade, based in part on the advancement of available technologies and results from repeat validation studies.

In recent history, diagnosis of the various forms of myeloid neoplasms has been based on a complex set of clinical, pathological, and biological criteria first introduced by the Polycythemia Vera Study Group (PVSG) in 1996<sup>[3, 4]</sup> and the World Health Organization (WHO) in 2001.<sup>[5]</sup> Both of these classifications use a combination of clinical, pathological, and/or biological criteria to arrive at a definitive diagnosis, predominantly reliant on status of Philadelphia chromosome presence. An important component of the diagnostic process is a clinical and laboratory assessment to rule out reactive or secondary causes of disease. Some diagnostic methods (e.g., bone marrow microscopy) are not well standardized and others (e.g., endogenous erythroid colony formation) are neither standardized nor widely available.<sup>[6-8]</sup> Diagnosis and monitoring of patients with Philadelphia chromosome negative myeloid neoplasms poses a challenge because many of the laboratory and clinical features of these diseases can be mimicked by other conditions such as reactive or secondary erythrocytosis, thrombocytosis or myeloid fibrosis. In addition, these entities can be difficult to distinguish on morphological bone marrow exam and diagnosis can be complicated by changing disease patterns.

The most up-to-date classification and benchmark for diagnosis of hematopoietic and lymphoid tissues is a result of collaboration between the Society for Hematopathology and the European Association for Haematopathology and is published by the WHO, most recently in 2022.<sup>[2]</sup> This edition varies from the previous versions with a refinement of diagnostic criteria and emphasis on actionable biomarkers. The current classification of myeloid neoplasm and acute leukemia subgroups are delineated in Table 2.

**Table 2. WHO Myeloid Proliferations and Neoplasms Classification, adapted<sup>[2]</sup>**

## **MYELOID NEOPLASMS, CHRONIC**

### **Myeloproliferative Neoplasms (MPN)**

Chronic myeloid leukemia (CML), *BCR-ABL* 1<sup>+</sup>  
 Polycythemia vera (PV)  
 Essential thrombocythemia (ET)  
 Primary myelofibrosis (PMF)  
 Chronic neutrophilic leukemia (CNL)  
 Chronic eosinophilic leukemia  
 Juvenile myelomonocytic leukemia (JMML)  
 MPN, not otherwise specified (NOS)

### **Mastocytosis**

Cutaneous mastocytosis  
 Systemic mastocytosis  
 Mast cell sarcoma

### **Myelodysplastic Neoplasms (MDS)**

MDS with low blasts and isolated 5q deletion  
 MDS with low blasts and SF3B1 mutation  
 MDS with low blasts, NOS  
 MDS with increased blasts  
 MDS with fibrosis  
 MDS, NOS  
 MDS with biallelic TP53 alteration (provisional)  
 MDS with other defined driver gene alterations

### **MDS / acute myeloid leukemia (MDS/AML)**

MDS/AML with *NMP1* mutation  
 MDS/AML with *MECOM* rearrangement  
 MDS/AML, NOS

### **MDS of childhood**

Refractory cytopenia of childhood  
 Childhood MDS

### **MDS with proliferative evolution**

Chronic myelomonocytic leukemia  
 MDS with proliferative evolution and neutrophilia  
 MDS with proliferative evolution, SF3B1 mutation and thrombocytosis  
 MDX with proliferative evolution, NOS

## **MYELOID NEOPLASMS, ACUTE**

### **AML with defining genetic abnormalities**

Acute promyelocytic leukemia  
 AML with *RUNX1::RUNX1T1* fusion  
 AML with *CBFB::MYH11* fusion  
 AML with *DEK::NUP214* fusion

AML with *RBM15::MRTFA* fusion

AML with *BCR::ABL1* fusion

AML with *NUP98* rearrangement

AML with *NPM1* mutation

AML with *NUP98* rearrangement

AML with other defined driver gene alterations

AML with myelodysplasia-related cytogenetics

### **AML, defined by differentiation**

AML with minimal differentiation

AML with without maturation

AML with maturation

Acute basophilic leukemia

AML with myelomonocytic differentiation

AML with monocytic differentiation

AML with plasmacytoid dendritic cell differentiation (provisional)

Pure erythroid leukemia

Acute megakaryoblastic leukemia

### **Myeloid sarcoma**

## **MYELOID NEOPLASMS, SECONDARY**

### **Myeloid neoplasms and proliferations associated with antecedent or predisposing conditions**

Myeloid neoplasm post cytotoxic therapy  
 Myeloid neoplasm associated with germline predisposition  
 AML following other hematolymphoid malignancy  
 Myeloid proliferations associated with Down syndrome  
 Myeloid neoplasm associated with malignant germ cell tumor

## **MYELOID/LYMPHOID NEOPLASMS AND OTHER LEUKAEMIAS OF AMBIGUOUS LINEAGE**

### **Myeloid/Lymphoid neoplasms with eosinophilia and defining gene rearrangement**

Myeloid/lymphoid neoplasm with *PDGFRA* rearrangement  
 Myeloid/lymphoid neoplasm with *PDGFRB* rearrangement  
 Myeloid/lymphoid neoplasm with *FGFR1* rearrangement  
 Myeloid/lymphoid neoplasms with *PMC1-JAK2* fusion

### **Acute leukemias of ambiguous lineage**

Mixed-phenotype acute leukaemia with *BCR-ABL1* fusion  
 Mixed-phenotype acute leukaemia with *KMT2A* rearrangement  
 Mixed-phenotype acute leukaemia, B/myeloid  
 Acute leukaemia of ambiguous lineage, NOS

It is important to note that the presence of any one or more of the gene variants included in this policy may not be sufficient to confirm a diagnosis, rather, testing may help support other clinical, laboratory, or pathological findings.

## TREATMENT MONITORING

CML represents one of the earliest examples of the use of molecular information to revolutionize patient management. A unique chromosomal change (the Philadelphia chromosome) and an accompanying unique gene rearrangement (*BCR-ABL*) resulting in a continuously activated tyrosine kinase enzyme were identified. These led to the development of a targeted tyrosine kinase inhibitor drug therapy (imatinib) that produces long-lasting remissions.

## REGULATORY STATUS

More than a dozen commercial laboratories currently offer a wide variety of diagnostic procedures for gene variant testing related to myeloid neoplasms and acute lymphoblastic leukemia. These tests are available as laboratory developed procedures under the U.S. Food and Drug Administration (FDA) enforcement discretion policy for laboratory developed tests (LDTs). Clinical laboratories may develop and validate tests in-house and market them as a laboratory service; LDTs must meet the general regulatory standards of Clinical Laboratory Improvement Act (CLIA) and laboratories that offer LDTs must be licensed by CLIA for high-complexity testing. To date, FDA does not require regulatory review of LDTs.

The FDA Centers for Devices and Radiological Health (CDRH), for Biologics Evaluation and Research (CBER), and for Drug Evaluation and Research (CDER) developed a draft guidance on in vitro companion diagnostic devices, which was released on July 14, 2011,<sup>[9]</sup> to address the “emergence of new technologies that can distinguish subsets of populations that respond differently to treatment.” As stated, the FDA encourages the development of treatments that depend on the use of companion diagnostic devices “when an appropriate scientific rationale supports such an approach.” In such cases, the FDA intends to review the safety and effectiveness of the companion diagnostic test as used with the therapeutic treatment that depends on its use. The rationale for co-review and approval is the desire to avoid exposing patients to preventable treatment risk.

The LeukoStrat<sup>®</sup> CDx *FLT3* Mutation Assay offered by Invivoscribe. According to Invivoscribe, the test is indicated at initial diagnosis of AML to determine eligibility for Rydapt<sup>®</sup> (midostaurin), Xospata<sup>®</sup> (gilteritinib), and Vanflyta<sup>®</sup> (quizartinib), and may also be used for risk stratification.<sup>[10]</sup> The assay includes internal tandem duplication variant testing for *FLT3* as well as variants in the tyrosine kinase domain. The assay is an FDA-approved companion diagnostic test for use with these medications and therefore may be standard of care in screening patients for use with this specific kinase inhibitor.

Abbott RealTime IDH2 is an *in vitro* polymerase chain reaction (PCR) assay for the qualitative detection of single nucleotide variants (SNVs) in the human isocitrate dehydrogenase-2 (IDH2) gene. The test aids in identifying acute myeloid leukemia patients for treatment with Idhifa<sup>®</sup> (enasidenib). Enasidenib is an oral medication used to treat patients with AML when the disease recurs after or does not respond to front-line therapies. The Abbott RealTime IDH2 assay received FDA premarket approval in August 2017.

## EVIDENCE SUMMARY

Human Genome Variation Society (HGVS) nomenclature is used to describe variants found in DNA and serves as an international standard.<sup>[11]</sup> It is being implemented for genetic testing medical evidence review updates starting in 2017. According to this nomenclature, the term “variant” is used to describe a change in a DNA or protein sequence, replacing previously-used terms, such as “mutation.” Pathogenic variants are variants associated with disease, while benign variants are not. The majority of genetic changes have unknown effects on human health, and these are referred to as variants of uncertain significance.

Validation of the clinical use of any genetic test focuses on three main principles:

1. The analytic validity of the test, which refers to the technical accuracy of the test in detecting a variant that is present or in excluding a variant that is absent;
2. The clinical validity of the test, which refers to the diagnostic performance of the test (sensitivity, specificity, positive and negative predictive values) in detecting clinical disease; and
3. The clinical utility of the test, i.e., how the results of the diagnostic test will be used to change management of the patient and whether these changes in management lead to clinically important improvements in health outcomes.

The focus of this review is on evidence related to the ability of test results to:

- Guide decisions in the clinical setting related to either treatment, management, or prevention, and
- Improve health outcomes as a result of those decisions.

## **BCR-ABL1 (*ABL1*) KINASE DOMAIN ANALYSIS**

Screening for *BCR-ABL1* kinase domain variants in chronic phase CML is recommended for patients with inadequate initial response to TKI treatment, those with evidence of loss of response, and for patients who have progressed to accelerated or blast phase CML.<sup>[3]</sup> The focus of the following discussion is on kinase domain point variants and treatment outcomes in systematic reviews.

In 2010, the Agency for Healthcare Research and Quality published a systematic review on *BCR-ABL1* pharmacogenetic testing for tyrosine kinase inhibitors in CML.<sup>[12]</sup> Thirty-one publications of *BCR-ABL1* testing met the eligibility criteria and were included in the review (20 of dasatinib, seven of imatinib, three of nilotinib, and one with various TKIs). The report concluded that the presence of any *BCR-ABL1* variant does not predict differential response to TKI therapy, although the presence of the T315I variant uniformly predicts TKI failure. However, during the public comment period the review was strongly criticized by respected pathology organizations for lack of attention to several issues that were subsequently insufficiently addressed in the final report. Importantly, the review grouped together studies that used kinase domain variant screening methods with those that used targeted methods and combined studies that used variant detection technologies with very different sensitivities. The authors dismissed the issues as related to analytic validity and beyond the scope of the report. However, in this clinical scenario assays with different intent (screening vs. targeted) and assays of very different sensitivities may lead to different clinical conclusions, so an understanding of these points is critical.

Branford (2009) summarized much of the available evidence regarding kinase domain variants detected at imatinib failure, and subsequent treatment success or failure with dasatinib or



nilotinib.<sup>[13]</sup> The T315I variant was most common; although about 100 variants have been reported, the seven most common (at residues T315, Y253, E255, M351, G250, F359, and H396) accounted for 60-66% of all variants. However, preexisting or emerging variants T315A, F317L/I/V/C, and V299L are associated with decreased clinical efficacy with dasatinib treatment following imatinib failure. Detection of the T315I variant at imatinib failure is associated with lack of subsequent response to high-dose imatinib, or to dasatinib or nilotinib. For these patients, allogeneic stem-cell transplantation remained the only available treatment until the advent of new agents such as ponatinib.<sup>[14]</sup> However these variants do not correspond to clinical significance, and based on clinical studies, the majority of imatinib-resistant variants remain sensitive to dasatinib and nilotinib.

Preexisting or emerging variants T315A, F317L/I/V/C, and V299L are associated with decreased clinical efficacy with dasatinib treatment following imatinib failure. Similarly, preexisting or emerging variants Y253H, E255K/V, and F359V/C have been reported for decreased clinical efficacy with nilotinib treatment following imatinib failure. In the survey reported by Branford, a total of 42% of patients tested had T315I or one of these dasatinib- or nilotinib-resistant variants. In the absence of any of these actionable variants, various treatment options are available. Note that these data have been obtained from studies in which patients were all initially treated with imatinib; no data are available regarding variants developing during first-line therapy with dasatinib or nilotinib.<sup>[15]</sup>

Unlike in CML, resistance in ALL to TKIs is less well studied. Resistance does not necessarily arise from dominant tumor clone(s), but possibly in response to TKI-driven selective pressure and/or by competition of other coexisting subclones.<sup>[16]</sup> In patients with ALL that are receiving a TKI, a rise in the BCR-ABL level while in hematologic complete response or clinical relapse warrants variant analysis.

## **ASXL1, CALR, IDH1, IDH2 AND TP53 IN MYELOID NEOPLASMS AND LEUKEMIA**

Testing for the *ASXL1*, *CALR*, *IDH1*, *IDH2* and *TP53* is required to meet WHO diagnostic criteria for patients with all of the most common Philadelphia-chromosome-negative MPNs.. The most recent revisions to the WHO criteria (2022) are heavily based on repeat validation studies.<sup>[2]</sup> The following evidence highlights the diagnostic and prognostic significance of *ASXL1*, *CALR*, *IDH1*, *IDH2* and *TP53* as specified by WHO diagnostic criteria and National Comprehensive Cancer Network (NCCN) guidelines.

### **ASXL1**

For chronic myelomonocytic leukemia (CMML), *ASXL1* is amongst the most frequently mutated genes, observed in 40-50% of CMML patients.<sup>[17-18]</sup> *ASXL1* is also reported to be associated with chromatin modification in MPNs, including polycythemia vera, as well as pre- and overt primary myelofibrosis.<sup>[19-20]</sup>

### **CALR**

Evidence for *CALR* demonstrates that a significant proportion of patients with myeloproliferative neoplasms and normal *JAK2* V617F status have a *CALR* variant.<sup>[21-23]</sup> Variants in exon 9 of *CALR* are found in 20-35% of all patients with ET and myelofibrosis. Fifty-two base pair deletions (*Type 1*) and five base pair insertions (*Type 2*) are the most common.

It is suggested that ET patients with *CALR* variants have lower polycythemic transformation rates, but not lower myelofibrotic transformation rate, compared with ET patients harboring a



*JAK2* variant. Chen (2014) reported a higher platelet count, younger age of diagnosis, lower leukocyte count, and decreased risk for thrombosis, compared with a *JAK2* positive ET population.<sup>[24]</sup> Tefferi (2014) reported survival and blast transformation in primary myelofibrosis (PMF) were significantly affected by variant status, though not in ET.<sup>[25]</sup> The outcome was best in *CALR*-variant patients and worst in *JAK2/CALR/MPL*-negative PMF patients. *CALR*-variant ET has also been associated with better thrombosis-free survival and lower leukocyte counts. However, overall survival has been reported as not different among *CALR*-variant and non-variant ET.<sup>[26 27]</sup>

## ***IDH1/2***

For PMF and ET, WHO criteria specify *IDH1/2* (as well as others, including *ASXL1*) as having diagnostic significance for those without *JAK2*, *CALR*, and *MPL* variants. In myeloproliferative neoplasms, *IDH1* and *IDH2* variants are among a growing number of higher-risk molecular markers. Both are associated with shorter overall survival and leukemia-free survival in patients with PMF and polycythemia vera.<sup>[20 28]</sup> In a study of the prognostic significance of *ASXL1*, *EZH2*, *SRSF2*, *IDH1* and *IDH2*, Vannucchi (2013) analyzed samples from 897 PMF patients (483 European patients and 396 from the Mayo clinical validation cohort). Median survival was significantly shorter (81 vs. 148 months,  $p < 0.0001$ ) in PMF patients with at least one of the genes.

## ***TP53***

Like *IDH1/2* described above, for PMF, *TP53* is associated with leukemic transformation, which is a common risk amongst patients with myeloproliferative neoplasms.<sup>[29]</sup> Furthermore, *TP53* is associated with inferior leukemia-free survival in those with ET. This progression is associated with poor clinical outcomes and resistance to standard AML therapies. Thus, *TP53* variants have also been analyzed to subdivide AML into prognostic subsets (see below). Additionally, *TP53* variants have been identified as one of the most common molecular abnormalities associated with myelodysplastic syndromes and may aid in diagnosis.<sup>[30-32]</sup>

## **ACUTE MYELOID LEUKEMIA**

AML is a group of diverse hematologic malignancies characterized by the clonal expansion of myeloid blasts in the bone marrow, blood, and/or other tissues. It is the most common type of leukemia in adults and is generally associated with a poor prognosis. It was estimated that in 2014, 18,860 people would be diagnosed with AML and 10,460 would die of the disease. Median age at diagnosis is 66 years, with approximately one in three patients diagnosed at 75 years of age or older.<sup>[33]</sup>

Conventional cytogenetic analysis (karyotyping) is a key component of the diagnostic evaluation of patients with suspected acute leukemia. The cytogenetic profile of the tumor is currently the most powerful predictor of prognosis in AML and is used to guide risk-adapted treatment strategies. Molecular variants, including those in *CEBPA*, *FLT3*, *KIT*, *NPM1*, *RUNX1*, and *TP53* genes, can be used to subdivide AML into prognostic subsets. (See Table 3.) Patients with better-prognosis disease based on cytogenetics (e.g., core-binding factor AML) who have a *c-KIT* variant in leukemic blast cells do just as poorly with post-remission standard chemotherapy as patients with cytogenetically poor-risk AML.<sup>[34]</sup> Similarly, individuals with cytogenetically normal AML (intermediate-prognosis disease) can be subcategorized into groups with better or worse prognosis based on the variant status of the *NPM1* and *FLT3* genes. Patients with variants in *NPM1* but without a *FLT3-ITD* fusion have post-remission

outcomes with standard chemotherapy that are similar to those with better-prognosis cytogenetics; in contrast, patients with any other combination of variants in those genes have outcomes similar to those with poor-prognosis cytogenetics.<sup>[35]</sup> A provisional category of AML with a *RUNX1* variant classifies de novo cases which are not associated with MDS-related cytogenetic abnormalities. This distinct group of AML patients also appears to have a worse prognosis than other AML types.<sup>[36-39]</sup>

The World Health Organization (WHO) classification of AML was adapted by the NCCN to estimate individual patient prognosis to guide management, as shown in Table 3:<sup>[40]</sup>

**Table 3. Risk Status of AML Based on Cytogenetic and Molecular Factors**

Risk Category	Genetic Abnormality
Favorable	t(8;21)(q22;q22.1)/ <i>RUNX1::RUNX1T1</i> inv(16)(p13.1;q22) or t(16;16)(p13.1;q22); <i>CBFB::MYH11</i> Mutated <i>NPM1</i> without <i>FLT3</i> -ITD bZIP in-frame mutated <i>CEBPA</i>
Intermediate	Mutated <i>NPM1</i> with <i>FLT3</i> -ITD Wild-type <i>NPM1</i> with <i>FLT3</i> -ITD (without adverse-risk genetic lesions) t(9;11)(p21.3;q23.3)/ <i>MLLT3::KMT2A</i> Cytogenetic and/or molecular abnormalities not classified as favorable or adverse
Poor/Adverse	t(6;9)(p23;q34.1)/ <i>DEK::NUP214</i> t(v;11q23.3)/ <i>KMT2A</i> -rearranged t(9;22)(q34.1;q11.2)/ <i>BCR::ABL1</i> t(8;16)(p11.2;p13.3)/ <i>KAT6A::CREBBP</i> inv(3)(q21.3;q26.2) or t(3;3)(q21.3;q26.2)/ <i>GATA2</i> , <i>MECOM(EVI1)</i> -5 or del(5q); -7; -17/abn(17p) Complex karyotype, monosomal karyotype Mutated <i>ASXL1</i> , <i>BCOR</i> , <i>EZH2</i> , <i>RUNX1</i> , <i>SF3B1</i> , <i>SRSF2</i> , <i>STAG2</i> , <i>U2AF1</i> , and/or <i>ZRSR2</i> Mutated <i>TP53</i>

## Genetic Testing for Molecular Subtypes of AML

A number of systematic reviews with meta-analyses have highlighted the evolving classification of AML into distinct molecular subtypes based on *CEBPA*, *FLT3-ITD*, *KIT*, *NPM1*, and *TP53*, particularly in patients with normal karyotype.<sup>[41-46]</sup> These studies support the WHO and NCCN risk status classifications, and additionally highlight the importance of *KIT* testing in the initial evaluation and for prognosis.

## PANEL TESTING FOR MYELOID NEOPLASMS

As indicated in NCCN guidelines and the WHO classification system, testing for variants in multiple genes may be indicated for diagnosis or treatment decisions in patients diagnosed with, or suspected of having, a myeloid neoplasm (see Practice Guideline Summary below). A number of studies have been published that describe the use of genetic panel tests that include these genes for diagnosis and prognosis of AML<sup>[47-51]</sup> and MDS<sup>[52-54]</sup>.

## PRACTICE GUIDELINE SUMMARY

### WORLD HEALTH ORGANIZATION

In 2016 the WHO published diagnostic criteria for myeloid neoplasms and acute leukemia, which include testing for a number of genetic variants, as shown in Table 2.<sup>[1]</sup> The 2022 major criteria for myeloproliferative neoplasms are unchanged.<sup>[2]</sup>

## NATIONAL COMPREHENSIVE CANCER NETWORK

The NCCN has published guidelines for Chronic Myeloid Leukemia (v.3.2025)<sup>[55]</sup>, Acute Lymphoblastic Leukemia (v.2.2025)<sup>[56]</sup>, which include recommendations regarding *BCR-ABL1* testing.

NCCN guidelines for Acute Myeloid Leukemia (v.2.2025)<sup>[40]</sup>, Myelodysplastic Syndromes (v.2.2025)<sup>[57]</sup>, and Myeloproliferative Neoplasms (v.1.2025)<sup>[58]</sup> include recommendations for testing a number of genes that have clinical significance for these disorders, including *JAK2*, *CALR*, *MPL*, *ASXL1*, *IDH1*, *IDH2*, *TP53*, *CEBPA*, *FLT3*, *KIT*, *NPM1*, and *RUNX1*.

## SUMMARY

### ***BCR/ABL1* (t(9;22)) TRANSLOCATION ANALYSIS, *JAK2*, *CALR*, *MPL*, *ASXL1*, *IDH1*, *IDH2*, *TP53*, *CEBPA*, *FLT3*, *KIT*, *NPM1* AND/OR *RUNX1***

There is enough research to show that *BCR/ABL1* (t(9;22)) translocation analysis (Philadelphia chromosome) and genetic testing for *JAK2*, *CALR*, *MPL*, *ASXL1*, *IDH1*, *IDH2*, *TP53*, *CEBPA*, *FLT3*, *KIT*, *NPM1* and/or *RUNX1* variants is important to guide diagnosis and treatment of myeloid neoplasms and leukemia. Additionally, these tests are recommended by clinical practice guidelines for various myeloid disorders. Therefore, testing for *BCR/ABL1* (t(9;22)) translocation analysis (Philadelphia chromosome) and genetic testing for *JAK2*, *CALR*, *MPL*, *ASXL1*, *IDH1*, *IDH2*, *TP53*, *CEBPA*, *FLT3*, *KIT*, *NPM1* and/or *RUNX1* variants is considered medically necessary for evaluation, diagnosis, and/or treatment monitoring for myeloid neoplasms and leukemia.

### **BCR-ABL KINASE DOMAIN (*ABL1*)**

In chronic myeloid leukemia, there is enough research to show clinical utility for evaluation of *ABL1* variants for tyrosine kinase inhibitor (TKI) resistance. TKI resistance in acute lymphoblastic leukemia (ALL) has not been studied as well as in CML. However, there is enough research to show *ABL1* genetic testing for evaluation of TKI resistance may lead to an improvement in health outcomes for patients with ALL who are receiving a TKI. Practice guidelines based on research recommend *ABL1* testing for ALL and CML in specific clinical scenarios. Therefore, *ABL1* genetic testing for evaluation of TKI resistance may be considered medically necessary when policy criteria are met. Due to insufficient evidence, evaluation of *ABL1* variants is considered investigational when policy criteria are not met.

### **TARGETED PANEL TESTING**

There is enough research to show that targeted panel testing may be important for diagnosis and guide treatment decisions for patients suspected of having or diagnosed with myeloproliferative neoplasms (MPN), myelodysplastic neoplasms (MDS), and myelodysplastic myeloproliferative neoplasms (MPN/MDS), including acute myeloid leukemia (AML). Clinical practice guidelines recommend panel testing for these disorders.

---

Therefore, targeted panel testing for MPN, MDS, MPN/MDS or AML may be considered medically necessary.

## NON-TARGETED PANEL TESTING

Non-targeted panels include testing for a large number of genes and are not targeted toward genes that have clinical significance for a specific type of hematolymphoid disorder. They often include testing for many genes that are not necessary to guide treatment, as well as testing for disorders that could be distinguished based on clinical presentation. There are no clinical practice guidelines based on research that recommend testing for all of the genes in these panels. Therefore, the use of non-targeted hematologic panel testing is considered investigational.

---

## REFERENCES

1. Arber DA, Orazi A, Hasserjian R, et al. The 2016 revision to the World Health Organization classification of myeloid neoplasms and acute leukemia. *Blood*. 2016;127(20):2391-405. PMID: 27069254
2. Khoury JD, Solary E, Abla O, et al. The 5th edition of the World Health Organization Classification of Haematolymphoid Tumours: Myeloid and Histiocytic/Dendritic Neoplasms. *Leukemia*. 2022;36(7):1703-19. PMID: 35732831
3. Murphy S, Peterson P, Iland H, et al. Experience of the Polycythemia Vera Study Group with essential thrombocythemia: a final report on diagnostic criteria, survival, and leukemic transition by treatment. *Semin Hematol*. 1997;34(1):29-39. PMID: 9025160
4. Pearson TC, Messinezy M. The diagnostic criteria of polycythaemia rubra vera. *Leuk Lymphoma*. 1996;22 Suppl 1:87-93. PMID: 8951778
5. Vardiman JW, Harris NL, Brunning RD. The World Health Organization (WHO) classification of the myeloid neoplasms. *Blood*. 2002;100(7):2292-302. PMID: 12239137
6. Tefferi A, Thiele J, Vardiman JW. The 2008 World Health Organization classification system for myeloproliferative neoplasms: order out of chaos. *Cancer*. 2009;115(17):3842-7. PMID: 19472396
7. Wilkins BS, Erber WN, Bareford D, et al. Bone marrow pathology in essential thrombocythemia: interobserver reliability and utility for identifying disease subtypes. *Blood*. 2008;111(1):60-70. PMID: 17885079
8. Baxter EJ, Scott LM, Campbell PJ, et al. Acquired mutation of the tyrosine kinase JAK2 in human myeloproliferative disorders. *Lancet*. 2005;365(9464):1054-61. PMID: 15781101
9. Chadalavada R, Lin E, Swafford V, et al. Comparative results of endoluminal gastroplasty and laparoscopic antireflux surgery for the treatment of GERD. *Surg Endosc*. 2004;18(2):261-5. PMID: 14691698
10. Invivoscribe. LeukoStrat® CDx FLT3 Mutation Assay. Secondary Invivoscribe. LeukoStrat® CDx FLT3 Mutation Assay [cited 3/27/2025]. 'Available from:' <https://invivoscribe.com/products/companion-diagnostics-cdx/>.
11. den Dunnen JT, Dalgleish R, Maglott DR, et al. HGVS Recommendations for the Description of Sequence Variants: 2016 Update. *Human mutation*. 2016;37(6):564-9. PMID: 26931183
12. Terasawa T, Dahabreh I, Castaldi PJ, et al. Systematic Reviews on Selected Pharmacogenetic Tests for Cancer Treatment: CYP2D6 for Tamoxifen in Breast

- Cancer, KRAS for anti-EGFR antibodies in Colorectal Cancer, and BCR-ABL1 for Tyrosine Kinase Inhibitors in Chronic Myeloid Leukemia [Internet]. Rockville (MD): Agency for Healthcare Research and Quality (US); 2010 Jun. 2010. PMID: 26065050
13. Branford S, Melo JV, Hughes TP. Selecting optimal second-line tyrosine kinase inhibitor therapy for chronic myeloid leukemia patients after imatinib failure: does the BCR-ABL mutation status really matter? *Blood*. 2009;114(27):5426-35. PMID: 19880502
  14. DiBaise JK, Brand RE, Quigley EM. Endoluminal delivery of radiofrequency energy to the gastroesophageal junction in uncomplicated GERD: efficacy and potential mechanism of action. *Am J Gastroenterol*. 2002;97(4):833-42. PMID: 12003416
  15. Alikian M, Gerrard G, Subramanian PG, et al. BCR-ABL1 kinase domain mutations: methodology and clinical evaluation. *American journal of hematology*. 2012;87(3):298-304. PMID: 22231203
  16. Fielding AK, Zakout GA. Treatment of Philadelphia chromosome-positive acute lymphoblastic leukemia. *Current hematologic malignancy reports*. 2013;8(2):98-108. PMID: 23475624
  17. Meggendorfer M, Roller A, Haferlach T, et al. SRSF2 mutations in 275 cases with chronic myelomonocytic leukemia (CMML). *Blood*. 2012;120(15):3080-8. PMID: 22919025
  18. Patnaik MM, Itzykson R, Lasho TL, et al. ASXL1 and SETBP1 mutations and their prognostic contribution in chronic myelomonocytic leukemia: a two-center study of 466 patients. *Leukemia*. 2014;28(11):2206-12. PMID: 24695057
  19. Guglielmelli P, Lasho TL, Rotunno G, et al. The number of prognostically detrimental mutations and prognosis in primary myelofibrosis: an international study of 797 patients. *Leukemia*. 2014;28(9):1804-10. PMID: 24549259
  20. Vannucchi AM, Lasho TL, Guglielmelli P, et al. Mutations and prognosis in primary myelofibrosis. *Leukemia*. 2013;27(9):1861-9. PMID: 23619563
  21. Tefferi A, Lasho TL, Finke CM, et al. CALR vs JAK2 vs MPL-mutated or triple-negative myelofibrosis: clinical, cytogenetic and molecular comparisons. *Leukemia*. 2014;28(7):1472-7. PMID: 24402162
  22. Klampfl T, Gisslinger H, Harutyunyan AS, et al. Somatic mutations of calreticulin in myeloproliferative neoplasms. *N Engl J Med*. 2013;369(25):2379-90. PMID: 24325356
  23. Rumi E, Pietra D, Pascutto C, et al. Clinical effect of driver mutations of JAK2, CALR, or MPL in primary myelofibrosis. *Blood*. 2014;124(7):1062-9. PMID: 24986690
  24. Chen CC, Gau JP, Chou HJ, et al. Frequencies, clinical characteristics, and outcome of somatic CALR mutations in JAK2-unmutated essential thrombocythemia. *Annals of hematology*. 2014;93(12):2029-36. PMID: 25015052
  25. Tefferi A, Guglielmelli P, Larson DR, et al. Long-term survival and blast transformation in molecularly annotated essential thrombocythemia, polycythemia vera, and myelofibrosis. *Blood*. 2014;124(16):2507-13; quiz 615. PMID: 25037629
  26. Ha JS, Kim YK. Calreticulin exon 9 mutations in myeloproliferative neoplasms. *Annals of laboratory medicine*. 2015;35(1):22-7. PMID: 25553276
  27. Yang Y, Wang X, Wang C, et al. A meta-analysis comparing clinical characteristics and outcomes in CALR-mutated and JAK2V617F essential thrombocythaemia. *International journal of hematology*. 2015;101(2):165-72. PMID: 25540065
  28. Tefferi A, Lasho TL, Guglielmelli P, et al. Targeted deep sequencing in polycythemia vera and essential thrombocythemia. *Blood advances*. 2016;1(1):21-30. PMID: 29296692

29. Rampal R, Ahn J, Abdel-Wahab O, et al. Genomic and functional analysis of leukemic transformation of myeloproliferative neoplasms. *Proceedings of the National Academy of Sciences of the United States of America*. 2014;111(50):E5401-10. PMID: 25516983
30. Haferlach T, Nagata Y, Grossmann V, et al. Landscape of genetic lesions in 944 patients with myelodysplastic syndromes. *Leukemia*. 2014;28(2):241-7. PMID: 24220272
31. Papaemmanuil E, Gerstung M, Malcovati L, et al. Clinical and biological implications of driver mutations in myelodysplastic syndromes. *Blood*. 2013;122(22):3616-27; quiz 99. PMID: 24030381
32. Bejar R, Stevenson K, Abdel-Wahab O, et al. Clinical effect of point mutations in myelodysplastic syndromes. *N Engl J Med*. 2011;364(26):2496-506. PMID: 21714648
33. Liersch R, Muller-Tidow C, Berdel WE, et al. Prognostic factors for acute myeloid leukaemia in adults--biological significance and clinical use. *Br J Haematol*. 2014;165(1):17-38. PMID: 24484469
34. Paschka P, Marcucci G, Ruppert AS, et al. Adverse prognostic significance of KIT mutations in adult acute myeloid leukemia with inv(16) and t(8;21): a Cancer and Leukemia Group B Study. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2006;24(24):3904-11. PMID: 16921041
35. Schlenk RF, Dohner K, Krauter J, et al. Mutations and treatment outcome in cytogenetically normal acute myeloid leukemia. *N Engl J Med*. 2008;358(18):1909-18. PMID: 18450602
36. Mendler JH, Maharry K, Radmacher MD, et al. RUNX1 mutations are associated with poor outcome in younger and older patients with cytogenetically normal acute myeloid leukemia and with distinct gene and MicroRNA expression signatures. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2012;30(25):3109-18. PMID: 22753902
37. Gaidzik VI, Bullinger L, Schlenk RF, et al. RUNX1 mutations in acute myeloid leukemia: results from a comprehensive genetic and clinical analysis from the AML study group. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2011;29(10):1364-72. PMID: 21343560
38. Schnittger S, Dicker F, Kern W, et al. RUNX1 mutations are frequent in de novo AML with noncomplex karyotype and confer an unfavorable prognosis. *Blood*. 2011;117(8):2348-57. PMID: 21148331
39. Tang JL, Hou HA, Chen CY, et al. AML1/RUNX1 mutations in 470 adult patients with de novo acute myeloid leukemia: prognostic implication and interaction with other gene alterations. *Blood*. 2009;114(26):5352-61. PMID: 19808697
40. National Comprehensive Cancer Network (NCCN) Guidelines. Acute Myeloid Leukemia. Secondary National Comprehensive Cancer Network (NCCN) Guidelines. Acute Myeloid Leukemia. [cited 3/27/2025]. 'Available from:' [https://www.nccn.org/professionals/physician\\_gls/pdf/aml.pdf](https://www.nccn.org/professionals/physician_gls/pdf/aml.pdf).
41. Li HY, Deng DH, Huang Y, et al. Favorable prognosis of biallelic CEBPA gene mutations in acute myeloid leukemia patients: a meta-analysis. *European journal of haematology*. 2015;94(5):439-48. PMID: 25227715
42. Wu X, Feng X, Zhao X, et al. Prognostic significance of FLT3-ITD in pediatric acute myeloid leukemia: a meta-analysis of cohort studies. *Molecular and cellular biochemistry*. 2016;420(1-2):121-8. PMID: 27435859
43. Port M, Bottcher M, Thol F, et al. Prognostic significance of FLT3 internal tandem duplication, nucleophosmin 1, and CEBPA gene mutations for acute myeloid leukemia



- patients with normal karyotype and younger than 60 years: a systematic review and meta-analysis. *Annals of hematology*. 2014;93(8):1279-86. PMID: 24801015
44. Ayatollahi H, Shajiei A, Sadeghian MH, et al. Prognostic Importance of C-KIT Mutations in Core Binding Factor Acute Myeloid Leukemia: A Systematic Review. *Hematology/oncology and stem cell therapy*. 2017;10(1):1-7. PMID: 27613372
  45. Chen W, Xie H, Wang H, et al. Prognostic Significance of KIT Mutations in Core-Binding Factor Acute Myeloid Leukemia: A Systematic Review and Meta-Analysis. *PloS one*. 2016;11(1):e0146614. PMID: 26771376
  46. Middeke JM, Herold S, Rucker-Braun E, et al. TP53 mutation in patients with high-risk acute myeloid leukaemia treated with allogeneic haematopoietic stem cell transplantation. *Br J Haematol*. 2016;172(6):914-22. PMID: 26771088
  47. Alonso CM, Llop M, Sargas C, et al. Clinical Utility of a Next-Generation Sequencing Panel for Acute Myeloid Leukemia Diagnostics. *J Mol Diagn*. 2019;21(2):228-40. PMID: 30576870
  48. Dunlap JB, Leonard J, Rosenberg M, et al. The combination of NPM1, DNMT3A, and IDH1/2 mutations leads to inferior overall survival in AML. *American journal of hematology*. 2019;94(8):913-20. PMID: 31145495
  49. Yu G, Yin C, Wu F, et al. Gene mutation profile and risk stratification in AML1-ETO-positive acute myeloid leukemia based on next-generation sequencing. *Oncol Rep*. 2019;42(6):2333-44. PMID: 31638252
  50. Salmoiraghi S, Cavagna R, Zanghì P, et al. High Throughput Molecular Characterization of Normal Karyotype Acute Myeloid Leukemia in the Context of the Prospective Trial 02/06 of the Northern Italy Leukemia Group (NILG). *Cancers (Basel)*. 2020;12(8). PMID: 32796597
  51. Chen X, Zhu H, Qiao C, et al. Next-generation sequencing reveals gene mutations landscape and clonal evolution in patients with acute myeloid leukemia. *Hematology*. 2021;26(1):111-22. PMID: 33491606
  52. Hamilton BK, Rybicki L, Hirsch C, et al. Mutation clonal burden and allogeneic hematopoietic cell transplantation outcomes in acute myeloid leukemia and myelodysplastic syndromes. *Bone Marrow Transplant*. 2019;54(8):1281-86. PMID: 30655603
  53. Zheng G, Chen P, Pallavajjalla A, et al. The diagnostic utility of targeted gene panel sequencing in discriminating etiologies of cytopenia. *American journal of hematology*. 2019;94(10):1141-48. PMID: 31350794
  54. Hunter AM, Komrokji RS, Yun S, et al. Baseline and serial molecular profiling predicts outcomes with hypomethylating agents in myelodysplastic syndromes. *Blood advances*. 2021;5(4):1017-28. PMID: 33591325
  55. National Comprehensive Cancer Network (NCCN) Guidelines. Chronic Myeloid Leukemia. Secondary National Comprehensive Cancer Network (NCCN) Guidelines. Chronic Myeloid Leukemia. [cited 3/27/2025]. 'Available from:' [https://www.nccn.org/professionals/physician\\_gls/pdf/cml.pdf](https://www.nccn.org/professionals/physician_gls/pdf/cml.pdf).
  56. National Comprehensive Cancer Network (NCCN) Guidelines. Acute Lymphoblastic Leukemia. Secondary National Comprehensive Cancer Network (NCCN) Guidelines. Acute Lymphoblastic Leukemia. [cited 3/27/2025]. 'Available from:' [https://www.nccn.org/professionals/physician\\_gls/pdf/all.pdf](https://www.nccn.org/professionals/physician_gls/pdf/all.pdf).
  57. National Comprehensive Cancer Network (NCCN) Guidelines. Myelodysplastic Syndromes. Secondary National Comprehensive Cancer Network (NCCN) Guidelines. Myelodysplastic Syndromes. [cited 3/27/2025]. 'Available from:' [https://www.nccn.org/professionals/physician\\_gls/pdf/mds.pdf](https://www.nccn.org/professionals/physician_gls/pdf/mds.pdf).



58. National Comprehensive Cancer Network (NCCN) Guidelines. Myeloproliferative Neoplasms. Secondary National Comprehensive Cancer Network (NCCN) Guidelines. Myeloproliferative Neoplasms. [cited 3/27/2025]. 'Available from:' [https://www.nccn.org/professionals/physician\\_gls/pdf/mpn.pdf](https://www.nccn.org/professionals/physician_gls/pdf/mpn.pdf).

## CODES

**NOTE:** BCR/ABL1 (t(9;22)) translocation analysis has specific CPT codes: 81206-8, 0016U, and 0040U. This differs from than BCR-ABL kinase domain (*ABL1*) variant analysis.

Codes	Number	Description
CPT	0016U	Oncology (hematolymphoid neoplasia), RNA, BCR/ABL1 major and minor breakpoint fusion transcripts, quantitative PCR amplification, blood or bone marrow, report of fusion not detected or detected with quantitation
	0017U	Oncology (hematolymphoid neoplasia), JAK2 mutation, DNA, PCR amplification of exons 12-14 and sequence analysis, blood or bone marrow, report of JAK2 mutation not detected or detected
	0023U	Oncology (acute myelogenous leukemia), DNA, genotyping of internal tandem duplication, p.D835, p.I836, using mononuclear cells, reported as detection or non-detection of FLT3 mutation and indication for or against the use of midostaurin
	0027U	JAK2 (Janus kinase 2) (eg, myeloproliferative disorder) gene analysis, targeted sequence analysis exons 12-15
	0040U	BCR/ABL1 (t(9;22)) (eg, chronic myelogenous leukemia) translocation analysis, major breakpoint, quantitative
	0046U	FLT3 (fms-related tyrosine kinase 3) (eg, acute myeloid leukemia) internal tandem duplication (ITD) variants, quantitative
	0049U	NPM1 (nucleophosmin) (eg, acute myeloid leukemia) gene analysis, quantitative
	0050U	Targeted genomic sequence analysis panel, acute myelogenous leukemia, DNA analysis, 194 genes, interrogation for sequence variants, copy number variants or rearrangements
	81120	IDH1 (isocitrate dehydrogenase 1 [NADP+], soluble) (eg, glioma), common variants (eg, R132H, R132C)
	81121	IDH2 (isocitrate dehydrogenase 2 [NADP+], mitochondrial) (eg, glioma), common variants (eg, R140W, R172M)
	81170	ABL1 (ABL proto-oncogene 1, non-receptor tyrosine kinase) (eg, acquired imatinib tyrosine kinase inhibitor resistance), gene analysis, variants in the kinase domain
	81175	ASXL1 (additional sex combs like 1, transcriptional regulator) (eg, myelodysplastic syndrome, myeloproliferative neoplasms, chronic myelomonocytic leukemia), gene analysis; full gene sequence
	81176	ASXL1 (additional sex combs like 1, transcriptional regulator) (eg, myelodysplastic syndrome, myeloproliferative neoplasms, chronic myelomonocytic leukemia), gene analysis; targeted sequence analysis (eg, EXON 12)
	81206	BCR/ABL1 (t(9;22)) (eg, chronic myelogenous leukemia) translocation analysis; major breakpoint, qualitative or quantitative
	81207	BCR/ABL1 (t(9;22)) (eg, chronic myelogenous leukemia) translocation analysis; minor breakpoint, qualitative or quantitative
	81208	BCR/ABL1 (t(9;22)) (eg, chronic myelogenous leukemia) translocation analysis; other breakpoint, qualitative or quantitative

Codes	Number	Description
	81218	<i>CEBPA</i> (CCAAT/enhancer binding protein [C/EBP], alpha) (eg, acute myeloid leukemia), gene analysis, full gene sequence
	81219	<i>CALR</i> (calreticulin) (eg, myeloproliferative disorders), gene analysis, common variants in exon 9
	81245	<i>FLT3</i> (fms-related tyrosine kinase 3) (eg, acute myeloid leukemia), gene analysis; internal tandem duplication (ITD) variants (ie, exons 14, 15)
	81246	<i>FLT3</i> (fms-related tyrosine kinase 3) (eg, acute myeloid leukemia), gene analysis; tyrosine kinase domain (TKD) variants (eg, D835, I836)
	81270	<i>JAK2</i> (Janus kinase 2) (eg, myeloproliferative disorder) gene analysis, p.Val617Phe (V617F) variant
	81272	<i>KIT</i> (v-kit Hardy-Zuckerman 4 feline sarcoma viral oncogene homolog) (eg, gastrointestinal stromal tumor [GIST], acute myeloid leukemia, melanoma), gene analysis, targeted sequence analysis (eg, exons 8, 11, 13, 17, 18)
	81273	<i>KIT</i> (v-kit Hardy-Zuckerman 4 feline sarcoma viral oncogene homolog) (eg, mastocytosis), gene analysis, D816 variant(s)
	81279	<i>JAK2</i> (Janus kinase 2) (eg, myeloproliferative disorder) targeted sequence analysis (eg, exons 12 and 13)
	81310	<i>NPM1</i> (nucleophosmin) (eg, acute myeloid leukemia) gene analysis, exon 12 variants
	81334	<i>RUNX1</i> (runt related transcription factor 1) (eg, acute myeloid leukemia, familial platelet disorder with associated myeloid malignancy), gene analysis, targeted sequence analysis (eg, EXONS 3-8)
	81338	<i>MPL</i> (MPL proto-oncogene, thrombopoietin receptor) (eg, myeloproliferative disorder) gene analysis; common variants (eg, W515A, W515K, W515L, W515R)
	81339	<i>MPL</i> (MPL proto-oncogene, thrombopoietin receptor) (eg, myeloproliferative disorder) gene analysis; sequence analysis, exon 10
	81351	<i>TP53</i> (tumor protein 53) (eg, Li-Fraumeni syndrome) gene analysis; full gene sequence
	81352	<i>TP53</i> (tumor protein 53) (eg, Li-Fraumeni syndrome) gene analysis; targeted sequence analysis (eg, 4 oncology)
	81353	<i>TP53</i> (tumor protein 53) (eg, Li-Fraumeni syndrome) gene analysis; known familial variant
	81401	Molecular pathology procedure, Level 2 - which includes <i>ABL1</i> ( <i>ABL proto oncogene 1, non-receptor tyrosine kinase</i> ) (eg, acquired imatinib resistance), T315I variant
	81402	Molecular pathology procedure, Level 3 (eg, >10 SNPs, 2-10 methylated variants, or 2-10 somatic variants [typically using non-sequencing target variant analysis], immunoglobulin and T-cell receptor gene rearrangements, duplication/deletion variants 1 exon)
	81403	Molecular pathology procedure, Level 4 (eg, analysis of single exon by DNA sequence analysis, analysis of >10 amplicons using multiplex PCR in 2 or more independent reactions, mutation scanning or duplication/deletion variants of 2-5 exons)
	81450	Hematolymphoid neoplasm or disorder, genomic sequence analysis panel, 5-50 genes, interrogation for sequence variants, and copy number variants or rearrangements, or isoform expression or mRNA expression levels, if performed; DNA analysis or combined DNA and RNA analysis
	81451	Hematolymphoid neoplasm or disorder, genomic sequence analysis panel, 5-50 genes, interrogation for sequence variants, and copy number variants or rearrangements, or isoform expression or mRNA expression levels, if performed; RNA analysis

<b>Codes</b>	<b>Number</b>	<b>Description</b>
	81455	Solid organ or hematolymphoid neoplasm or disorder, 51 or greater genes, genomic sequence analysis panel, interrogation for sequence variants and copy number variants or rearrangements, or isoform expression or mRNA expression levels, if performed DNA analysis or combined DNA and RNA analysis
	81456	Solid organ or hematolymphoid neoplasm or disorder, 51 or greater genes, genomic sequence analysis panel, interrogation for sequence variants and copy number variants or rearrangements, or isoform expression or mRNA expression levels, if performed; RNA analysis
HCPCS	None	

**Date of Origin:** August 2010